arXiv:gr-qc/0601008v2 9 Aug 2007

Quantum Field Theory in Curved Spacetime*

Bernard S. Kay
Department of Mathematics, University of York, York YO10 5DD, U.K.

February 3, 2008

1 Introduction and Preliminaries

Quantum Field Theory (QFT) in Curved Spacetime is a hybrid approximate theory in which quantum matter fields are assumed to propagate in a fixed classical background gravitational field. Its basic physical prediction is that strong gravitational fields can polarize the vacuum and, when time-dependent, lead to pair creation just as a strong and/or time-dependent electromagnetic field can polarize the vacuum and/or give rise to pair-creation of charged particles. One expects it to be a good approximation to full quantum gravity provided the typical frequencies of the gravitational background are very much less than the Planck frequency $(c^5/G\hbar)^{1/2} \sim 10^{43}s^{-1}$) and provided, with a suitable measure for energy, the energy of created particles is very much less than the energy of the background gravitational field or of its matter sources. Undoubtedly the most important prediction of the theory is the Hawking effect, according to which a, say spherically symmetric, classical black hole of mass M will emit thermal radiation at the Hawking temperature $T = (8\pi M)^{-1}$ (here and from now on, we use Planck units where G, c, $\hbar$ and k [Boltzmann's constant] are all taken to be 1.)

On the mathematical side, the need to formulate the laws and derive the general properties of QFT on non-flat spacetimes forces one to state and prove results in local terms and, as a byproduct, thereby leads to an improved perspective on flat-space-time QFT too. It's also interesting to formulate QFT on idealized spacetimes with particular global geometrical features. Thus, QFT on spacetimes with bifurcate Killing horizons is intimately related to the Hawking effect; QFT on spacetimes with closed timelike curves is intimately related to the question whether the laws of physics permit the manufacture of a time-machine.

*Published as *Encyclopedia of Mathematical Physics*, J.-P. Françoise, G. Naber and T.S. Tsou, eds., Academic (Elsevier), Amsterdam, New York, London 2006, Vol. 4, pp 202-212

As is standard in General Relativity, a curved spacetime is modelled mathematically as a (paracompact, Hausdorff) manifold $\mathcal{M}$ equipped with a pseudo-Riemannian metric g of signature $(-,+++)$ (we follow the conventions of the standard text "*Gravitation*" by Misner Thorne and Wheeler [W.H. Freeman, San Francisco, 1973]). We shall also assume, except where otherwise stated, our spacetime to be *globally hyperbolic*, i.e. that $\mathcal{M}$ admits a *global time coordinate*, by which we mean a global coordinate t such that each constant-t surface is a smooth Cauchy surface i.e. a smooth spacelike 3-surface cut exactly once by each inextendible causal curve. (Without this default assumption, extra problems arise for QFT which we shall briefly mention in connection with the time-machine question in Section 6.) In view of this definition, globally hyperbolic spacetimes are clearly time-orientable and we shall assume a choice of time-orientation has been made so we can talk about the "future" and "past" directions. Modern formulations of the subject take, as the fundamental mathematical structure modelling the quantum field, a $*$-algebra $\mathcal{A}$ (with identity I) together with a family of local sub $*$-algebras $\mathcal{A}(\mathcal{O})$ labelled by bounded open regions $\mathcal{O}$ of the spacetime $(\mathcal{M}, g)$ and satisfying the *isotony* or *net* condition that $\mathcal{O}_1 \subset \mathcal{O}_2$ implies $\mathcal{A}(\mathcal{O}_1)$ is a subalgebra of $\mathcal{A}(\mathcal{O}_2)$, as well as the condition that whenever $\mathcal{O}_1$ and $\mathcal{O}_2$ are spacelike separated, then $\mathcal{A}(\mathcal{O}_1)$ and $\mathcal{A}(\mathcal{O}_2)$ commute.

Standard concepts and techniques from algebraic quantum theory are then applicable: In particular, *states* are defined to be positive (this means $\omega(A^*A) \geq 0 \ \forall A \in \mathcal{A}$) normalized (this means $\omega(I) = 1$) linear functionals on $\mathcal{A}$. One distinguishes between *pure* states and *mixed* states, only the latter being writeable as non-trivial convex combinations of other states. To each state, ω, the *GNS-construction* associates a representation, ρ_ω, of $\mathcal{A}$ on a Hilbert space $\mathcal{H}_\omega$ together with a cyclic vector $\Omega \in \mathcal{H}_\omega$ such that

$$\omega(A) = \langle \Omega | \rho_\omega(A) \Omega \rangle$$

(and the *GNS triple* $(\rho_\omega, \mathcal{H}, \Omega)$ is unique up to equivalence). There are often technical advantages in formulating things so that the $*$-algebra is a C*-algebra. Then the GNS representation is as everywhere-defined bounded operators and is irreducible if and only if the state is pure. A useful concept, due to Haag, is the *folium* of a given state ω which may be defined to be the set of all states ω_σ which arise in the form $\mathrm{Tr}(\sigma \rho_\omega(\cdot))$ where σ ranges over the density operators (trace-class operators with unit trace) on $\mathcal{H}_\omega$.

Given a state, ω, and an automorphism, α, which preserves the state (i.e. $\omega \circ \alpha = \omega$) then there will be a unitary operator, U, on H_ω which *implements* α in the sense that $\rho_\omega(\alpha(A)) = U^{-1} \rho_\omega(A) U$ and U is chosen uniquely by the condition $U\Omega = \Omega$.

On a *stationary* spacetime, i.e. one which admits a one-parameter group of isometries whose integral curves are everywhere timelike, the algebra will inherit a one-parameter group (i.e. satisfying $\alpha(t_1) \circ \alpha(t_2) = \alpha(t_1 + t_2)$) of

time-translation automorphisms, $\alpha(t)$, and, given any stationary state (i.e. one which satisfies $\omega \circ \alpha(t) = \omega \quad \forall t \in \mathbf{R}$) these will be implemented by a one-parameter group of unitaries, $U(t)$, on its GNS Hilbert space satsifying $U(t)\Omega = \Omega$. If $U(t)$ is strongly continuous so that it takes the form e^{-iHt} and if the Hamiltonian, H, is positive, then ω is said to be a *ground state*. Typically one expects ground states to exist and often be unique.

Another important class of stationary states for the algebra of a stationary spacetime is the class of *KMS states*, ω^β, at inverse temperature β; these have the physical interpretation of thermal equilibrium states. In the GNS representation of one of these, the automorphisms are also implemented by a strongly-continuous unitary group, e^{-iHt}, which preserves Ω but (in place of H positive) there is a complex conjugation, J, on H_ω such that

$$e^{-\beta H/2}\rho_\omega(A)\Omega = J\rho_\omega(A^*)\Omega \tag{1}$$

for all $A \in \mathcal{A}$. An attractive feature of the subject is that its main qualitative features are already present for linear field theories and, unusually in comparison with other questions in QFT, these are susceptible of a straightforward explicit and rigorous mathematical formulation. In fact, as our principal example, we give, in Section 2 a construction for the field algebra for the quantized real linear Klein Gordon equation

$$(\Box_g - m^2 - V)\phi = 0 \tag{2}$$

of mass m on a globally hyperbolic spacetime $(\mathcal{M}, g)$. Here, $\Box_g$ denotes the Laplace Beltrami operator $g^{ab}\nabla_a\partial_b$ $(= (|\det(g)|)^{-1/2}\partial_a((|\det(g)|^{1/2}g^{ab}\partial_b))$. We include a scalar external background classical field, V in addition to the external gravitational field represented by g. In case m is zero, taking V to equal $R/6$ where R denotes the Riemann scalar, makes the equation *conformally invariant*.

The main new feature of quantum field theory in curved spacetime (present already for linear field theories) is that, in a general (neither flat, nor stationary) spacetime there will not be any single preferred state but rather a family of preferred states, members of which are best regarded as on an equal footing with one-another. It is this feature which makes the above algebraic framework particularly suitable, indeed essential, to a clear formulation of the subject. Conceptually it is this feature which takes the most getting used to. In particular, one must realize that, as we shall explain in Section 3, the interpretation of a state as having a particular "particle-content" is in general problematic because it can only be relative to a particular choice of "vacuum" state and, depending on the spacetime of interest, there may be one state or several states or, frequently, no states at all which deserve the name "vacuum" and even when there are states which deserve this name, they will often only be defined in some approximate or asymptotic or transient sense or only on some subregion of the spacetime.

Concomitantly, one does not expect global observables such as the "particle number" or the quantum Hamiltonian of flat-spacetime free field theory to generalize to a curved spacetime context, and for this reason local observables play a central role in the theory. The quantized stress-energy tensor is a particularly natural and important such local observable and the theory of this is central to the whole subject. A brief introduction to it is given in Section 4.

This is followed by a further Section 5 on the Hawking and Unruh effects and a brief Section 6 on the problems of extending the theory beyond the "default" setting, to non-globally hyperbolic spacetimes. Finally, Section 7 briefly mentions a number of other interesting and active areas of the subject as well as issuing a few *warnings* to be borne in mind when reading the literature.

2 Construction of $*$-Algebra(s) for a Real Linear Scalar Field on Globally Hyperbolic Spacetimes and Some General Theorems

On a globally hyperbolic spacetime, the classical equation (2) admits well-defined *advanced* and *retarded Green* functions (strictly bidistributions) Δ^A and Δ^R and the standard covariant quantum free real (or "Hermitian") scalar field commutation relations familiar from Minkowski spacetime free field theory naturally generalize to the (heuristic) equation

$$[\hat{\phi}(x), \hat{\phi}(y)] = i\Delta(x,y)I$$

where Δ is the *Lichnérowicz commutator function* $\Delta = \Delta^A - \Delta^R$. Here, the " $\hat{}$ " on the quantum field $\hat{\phi}$ serves to distinguish it from a classical solution ϕ. In mathematical work, one does not assign a meaning to the field at a point itself, but rather aims to assign meaning to *smeared fields* $\hat{\phi}(F)$ for all real-valued test functions $F \in C_0^\infty(\mathcal{M})$ which are then to be interpreted as standing for $\int_{\mathcal{M}} \hat{\phi}(x) f(x) |\det(g)|^{\frac{1}{2}} d^4x$. In fact, it is straightforward to define a *minimal field algebra* (see below) $\mathcal{A}_{\mathrm{min}}$ generated by such $\hat{\phi}(F)$ which satisfy the suitably smeared version

$$[\hat{\phi}(F), \hat{\phi}(G)] = i\Delta(F,G)I$$

of the above commutation relations together with Hermiticity (i.e. $\hat{\phi}(F)^* = \hat{\phi}(F)$), the property of being a weak solution of the equation (2) (i.e. $\hat{\phi}((\Box_g - m^2 - V)F) = 0 \quad \forall F \in C_0^\infty(\mathcal{M})$) and linearity in test functions. There is a technically different alternative formulation of this minimal algebra, which is known as the *Weyl algebra*, which is constructed to be the C* algebra generated by operators $W(F)$ (to be interpreted as standing for

$\exp(i\int_{\mathcal{M}}\hat{\phi}(x)f(x)|\det(g)|^{\frac{1}{2}}d^4x)$ satisfying

$$W(F_1)W(F_2) = \exp(-i\Delta(F_1,F_2)/2)W(F_1+F_2)$$

together with $W(F)^* = W(-F)$ and $W((\Box_g - m^2 - V)F) = I$. With either the minimal algebra or the Weyl algebra one can define, for each bounded open region $\mathcal{O}$, subalgebras $A(\mathcal{O})$ as generated by the $\hat{\phi}(\cdot)$ (or the $W(\cdot)$) smeared with test functions suppported in $\mathcal{O}$ and verify that they satisfy the above "net" condition and commutativity at spacelike separation.

Specifying a state, ω, on $\mathcal{A}_{\min}$ is tantamount to specifying its collection of n-point distributions (i.e. smeared n-point functions) $\omega(\hat{\phi}(F_1)\dots\hat{\phi}(F_n))$ (In the case of the Weyl algebra, one restricts attention to "regular" states for which the map $F \to \omega(W(F))$ is sufficiently often differentiable on finite dimensional subspaces of $C_0^\infty(\mathcal{M})$ and defines the n-point distributions in terms of derivatives with respect to suitable parameters of expectation values of suitable Weyl algebra elements.) A particular role is played in the theory by the *quasi-free* states for which all the *truncated* n-point distributions except for $n = 2$ vanish. Thus all the n-point distributions for odd n vanish while the 4-point distribution is made out of the 2-point distribution according to

$$\omega(\hat{\phi}(F_1)\hat{\phi}(F_2)\hat{\phi}(F_3)\hat{\phi}(F_4)) = \omega(\hat{\phi}(F_1)\hat{\phi}(F_2))\omega(\hat{\phi}(F_3)\hat{\phi}(F_4))$$

$$+\omega(\hat{\phi}(F_1)\hat{\phi}(F_3))\omega(\hat{\phi}(F_2)\hat{\phi}(F_4)) + \omega(\hat{\phi}(F_1)\hat{\phi}(F_4))\omega(\hat{\phi}(F_2)\hat{\phi}(F_3))$$

etc. The anticommutator distribution

$$G(F_1,F_2) = \omega(\hat{\phi}(F_1)\hat{\phi}(F_2)) + \omega(\hat{\phi}(F_2)\hat{\phi}(F_1)) \tag{3}$$

of a quasi-free state (or indeed of any state) will satisfy the properties (for all test functions F, F_1, F_2 etc.):

(a) *(symmetry)* $G(F_1,F_2) = G(F_2,F_1)$

(b) *(weak bisolution property)*

$$G((\Box_g - m^2 - V)F_1, F_2) = 0 = G(F_1, (\Box_g - m^2 - V)F_2)$$

(c) *(positivity)* $G(F,F) \geq 0$ and $G(F_1,F_1)^{1/2}G(F_2,F_2)^{1/2} \geq |\Delta(F_1,F_2)|$

and it can be shown that, to every bilinear functional G on $C_0^\infty(\mathcal{M})$ satisfying (a), (b) and (c), there is a quasi-free state with two-point distribution $(1/2)(G+i\Delta)$. One further declares a quasi-free state to be *physically admissible* only if (for pairs of points in sufficiently small convex neighbourhoods)

(d) *(Hadamard condition)* "$G(x_1,x_2) = \frac{1}{2\pi^2}\left(u(x_1,x_2)\mathrm{P}\frac{1}{\sigma} + v(x_1,x_2)\log|\sigma| + w(x_1,x_2)\right)$"

This last condition expresses the requirement that (locally) the two-point distribution actually "is" (in the usual sense in which one says that a distribution "is" a function) a smooth function for pairs of non-null-separated points, and at the same time requires that the two-point distribution be singular at pairs of null-separated points and locally specifies the nature of the singularity for such pairs of points with a leading "principal part of $1/\sigma$" type singularity and a subleading "$\log|\sigma|$" singularity where σ denotes the square of the geodesic distance between x_1 and x_2. u (which satisfies $u(x_1, x_2) = 1$ when $x_1 = x_2$) and v are certain smooth two-point functions determined in terms of the local geometry and the local values of V by something called the *Hadamard procedure* while the smooth two-point function w depends on the state. We shall omit the details. The important point is that this Hadamard condition on the two-point distribution is believed to be the correct generalization to a curved-spacetime of the well-known universal short-distance behaviour shared by the truncated two-point distributions of all physically relevant states for the special case of our theory when the spacetime is flat (and V vanishes). In the latter case, u reduces to 1, and v to a simple power series $\sum_{n=0}^{\infty} v_n \sigma^n$ with $v_0 = m^2/4$ etc.

Actually, it is known (this is the content of "Kay's Conjecture" which was proved by M. Radzikowski in 1992) that (a), (b), (c) and (d) together imply that the two-point distribution is nonsingular at all pairs of (not necessarily close-together) spacelike-separated points. More important than this result itself is a reformulation of the Hadamard condition in terms of the concepts of *microlocal analysis* which Radzikowski originally introduced as a tool towards its proof.

(d') *(Wave Front Set [or Microlocal] Spectrum Condition)* $\mathrm{WF}(G + i\Delta) = \{(x_1, p_1; x_2, p_2) \in T^*(\mathcal{M}\times\mathcal{M})\setminus\mathbf{0} | x_1$ and x_2 lie on a single null geodesic, p_1 is tangent to that null geodesic and future pointing, and p_2 when parallel transported along that null geodesic from x_2 to x_1 equals $-p_1\}$

For the gist of what this means, it suffices to know that to say that an element (x, p) of the cotangent bundle of a manifold (excluding the zero section $\mathbf{0}$) is in the *wave front set*, WF, of a given distribution on that manifold may be expressed informally by saying that that distribution is singular *at* the point x *in the direction* p. (And here the notion is applied to $G + i\Delta$, thought of as a distribution on the manifold $\mathcal{M} \times \mathcal{M}$.)

We remark that generically (and, e.g., always if the spatial sections are compact and $m^2+V(x)$ is everywhere positive) the Weyl algebra for equation (2) on a given stationary spacetime will have a unique ground state and unique KMS states at each temperature and these will be quasi-free and Hadamard.

Quasi-free states are important also because of a theorem of R. Verch (1994, in verification of another conjecture of Kay) that (in the Weyl algebra

framework) on the algebra of any bounded open region, the folia of the quasi-free Hadamard states coincide. With this result one can extend the notion of physical admissibility to not-necessarily-quasi-free states by demanding that, to be admissible, a state belong to the resulting common folium when restricted to the algebra of each bounded open region; equivalently that it be a *locally normal* state on the resulting natural extension of the net of local Weyl algebras to a net of local W^* *algebras*.

3 Particle Creation and the Limitations of the Particle Concept

Global hyperbolicity also entails that the Cauchy problem is well posed for the classical field equation (2) in the sense that for every Cauchy surface, $\mathcal{C}$, and every pair (f, p) of Cauchy data in $C_0^\infty(\mathcal{C})$, there exists a unique solution ϕ in $C_0^\infty(\mathcal{M})$ such that $f = \phi|_\mathcal{C}$ and $p = |\det(g)|^{1/2} g^{0b}\partial_b\phi|_\mathcal{C}$. Moreover ϕ has compact support on all other Cauchy surfaces. Given a global time coordinate t, increasing towards the future, foliating $\mathcal{M}$ into a family of contant-t Cauchy surfaces, $\mathcal{C}_t$, and given a choice of global time-like vector field τ^a (for example, $\tau^a = g^{ab}\partial_b t$) enabling one to identify all the $\mathcal{C}_t$, say with $\mathcal{C}_0$, by identifying points cut by the same integral curve of τ^a, a single such classical solution ϕ may be pictured as a family $\{(f_t, p_t) : t \in \mathbf{R}\}$ of time-evolving Cauchy data on $\mathcal{C}_0$. Moreover, since (2) implies, for each pair of classical solutions, ϕ_1, ϕ_2, the conservation (i.e. $\partial_a j^a = 0$) of the current $j^a = |\det(g)|^{1/2} g^{ab}(\phi_1\partial_b\phi_2 - \phi_2\partial_b\phi_1)$, the symplectic form (on $C_0^\infty(\mathcal{C}_0) \times C_0^\infty(\mathcal{C}_0)$)

$$\sigma((f_t^1, p_t^1); (f_t^2, p_t^2)) = \int_{\mathcal{C}_0} (f_t^1 p_t^2 - p_t^1 f_t^2) d^3x$$

will be conserved in time. (d^3x denotes $dx^1 \wedge dx^2 \wedge dx^3$.)

Corresponding to this picture of classical dynamics, one expects there to be a description of quantum dynamics in terms of a family of sharp-time quantum fields (φ_t, π_t) on $\mathcal{C}_0$, satisfying heuristic canonical commutation relations

$$[\varphi_t(\mathbf{x}), \varphi_t(\mathbf{y})] = 0, \quad [\pi_t(\mathbf{x}), \pi_t(\mathbf{y})] = 0, \quad [\varphi_t(\mathbf{x}), \pi_t(\mathbf{y})] = i\delta^3(\mathbf{x}, \mathbf{y})I$$

and evolving in time according to the same dynamics as the Cauchy data of a classical solution. (Both these expectations are correct because the field equation is linear.) An elegant way to make rigorous mathematical sense of these expectations is in terms of a $*$-algebra with identity generated by Hermitian objects "$\sigma((\varphi_0, \pi_0); (f, p))$" ("symplectically smeared sharp-time fields at $t = 0$") satisfying linearity in f and p together with the commutation relations

$$[\sigma((\varphi_0, \pi_0); (f^1, p^1)), \sigma((\varphi_0, \pi_0); (f^2, p^2))] = i\sigma((f^1, p^1); (f^2, p^2))I$$

and to define (symplectically smeared) time-t sharp-time fields by demanding

$$\sigma((\varphi_t, \pi_t); (f_t, p_t)) = \sigma((\varphi_0, \pi_0); (f_0, p_0))$$

where (f_t, p_t) is the classical time-evolute of (f_0, p_0). This $*$-algebra of sharp-time fields may be identified with the (minimal) field $*$-algebra of the previous section, the $\hat{\phi}(F)$ of the previous section being identified with $\sigma((\varphi_0, \pi_0); (f, p))$ where (f, p) are the Cauchy data at $t = 0$ of $\Delta * F$. (This identification is of course many-one since $\hat{\phi}(F) = 0$ whenever F arises as $(\Box_g - m^2 - V)G$ for some test function $G \in C_0^\infty(\mathcal{M})$.)

Specializing momentarily to the case of the free scalar field $(\Box - m^2)\phi = 0$ $(m \neq 0)$ in Minkowski space with a flat $t = 0$ Cauchy surface, the "symplectically smeared" two-point function of the usual ground state ("Minkowski vacuum state"), ω_0, is given, in this formalism, by

$$\omega_0(\sigma((\varphi, \pi); (f^1, p^1))\sigma((\varphi, \pi); (f^2, p^2))) = \frac{1}{2}(\langle f^1 | \mu f^2 \rangle + \langle p^1 | \mu^{-1} p^2 \rangle + i\sigma((f^1, p^1); (f^2, p^2)) \tag{4}$$

where the inner products are in the *one-particle Hilbert space* $\mathcal{H} = L^2_{\mathbf{C}}(\mathbf{R}^3)$ and $\mu = (m^2 - \nabla^2)^{1/2}$. The GNS representation of this state may be concretely realised on the familiar *Fock space* $\mathcal{F}(\mathcal{H})$ over $\mathcal{H}$ by

$$\rho_0(\sigma((\varphi, \pi); (f, p))) = -i(\hat{a}^\dagger(\mathrm{a}) - (\hat{a}^\dagger(\mathrm{a}))^*)$$

where a denotes the element of $\mathcal{H}$

$$\mathrm{a} = \frac{(\mu^{1/2} f + i\mu^{-1/2} p)}{\sqrt{2}}$$

(we note in passing that, if we equip $\mathcal{H}$ with the symplectic form $2\mathrm{Im}\langle \cdot | \cdot \rangle$, then $K : (f, p) \mapsto \mathrm{a}$ is a symplectic map) and $\hat{a}^\dagger(\mathrm{a})$ is the usual smeared *creation operator* ($=$"$\int \hat{a}^\dagger(\mathbf{x})\mathrm{a}(\mathbf{x})d^3\mathbf{x}$") on $\mathcal{F}(\mathcal{H})$ satisfying

$$[(\hat{a}^\dagger(\mathrm{a}^1))^*, \hat{a}^\dagger(\mathrm{a}^2)] = \langle \mathrm{a}^1 | \mathrm{a}^2 \rangle_{\mathcal{H}} I.$$

The usual (smeared) *annihilation operator*, $\hat{a}(\mathrm{a})$, is $(\hat{a}^\dagger(C\mathrm{a}))^*$ where C is the natural complex conjugation, $\mathrm{a} \mapsto \mathrm{a}^*$ on $\mathcal{H}$. Both of these operators annihilate the *Fock vacuum vector* $\Omega^{\mathcal{F}}$. In this representation, the one parameter group of time-translation automorphisms

$$\alpha(t) : \sigma((\varphi_0, \pi_0); (f, p)) \mapsto \sigma((\varphi_t, \pi_t); (f, p)) \tag{5}$$

is implemented by $\exp(-iHt)$ where H is the second quantization of μ (i.e. the operator otherwise known as $\int \mu(k)\hat{a}^\dagger(k)\hat{a}(k)d^3k$) on $\mathcal{F}(\mathcal{H})$.

The most straightforward (albeit physically artificial) situation involving "particle creation" in a curved spacetime concerns a globally hyperbolic spacetime which, outside of a compact region, is isometric to Minkowski

space with a compact region removed – i.e. to a globally hyperbolic spacetime which is flat except inside a localized "bump" of curvature. See Figure 1. (One could also allow the function V in (2) to be non-zero inside the bump.) On the field algebra (defined as in the previous section) of such a spacetime, there will be an "in" vacuum state (which may be identified with the Minkowski vacuum to the past of the bump) and an "out" vacuum state (which may be identified with the Minkowski vacuum to the future of the bump) and one expects e.g. the "in vacuum" to arise as a many particle state in the GNS representation of the "out vacuum" corresponding to the creation of particles out of the vacuum by the bump of curvature.



Figure 1 A spacetime which is flat outside of a compact bump of curvature.

In the formalism of this section, if we choose our global time coordinate on such a spacetime so that, say, the $t = 0$ surface is to the past of the bump and the $t = T$ surface to its future, then the single automorphism $\alpha(T)$ (defined as in (5)) encodes the overall effect of the bump of curvature on the quantum field and one can ask whether it is implemented by a unitary operator in the GNS representation of the Minkowski vacuum state (4).

This question may be answered by referring to the real-linear map $\mathcal{T} : \mathcal{H} \to \mathcal{H}$ which sends $a_T = 2^{-1/2}(\mu^{1/2} f_T + i\mu^{-1/2} p_T)$ to $a_0 = 2^{-1/2}(\mu^{1/2} f_0 + i\mu^{-1/2} p_0)$. By the conservation in time of σ and the symplecticity, noted in passing above, of the map $K : (f, p) \mapsto a$, this satisfies the defining relation

$$\mathrm{Im}\langle \mathcal{T}\mathrm{a}^1 | \mathcal{T}\mathrm{a}^2 \rangle = \mathrm{Im}\langle \mathrm{a}^1 | \mathrm{a}^2 \rangle$$

of a *classical Bogoliubov transformation*. Splitting $\mathcal{T}$ into its complex-linear and complex-antilinear parts by writing

$$\mathcal{T} = \alpha + \beta C$$

where α and β are complex linear operators, this relation may alternatively be expressed in terms of the pair of relations

$$\alpha^* \alpha - \bar{\beta}^* \bar{\beta} = I, \quad \bar{\alpha}^* \bar{\beta} = \beta^* \alpha$$

where $\bar{\alpha} = C\alpha C$, $\bar{\beta} = C\beta C$.

We remark that there is an easy-to-visualize equivalent way of defining α and β in terms of the analysis, to the past of the bump, into *positive and negative-frequency parts* of complex solutions to (2) which are purely *positive-frequency* to the future of the bump. In fact, if, for any element $a \in \mathcal{H}$, we identify the positive frequency solution to the Minkowski-space Klein Gordon equation

$$\phi^{\mathrm{pos}}_{\mathrm{out}}(t, \mathbf{x}) = ((2\mu)^{-1/2} \exp(-i\mu t)a)(\mathbf{x})$$

with a complex solution to (2) to the future of the bump, then (it may easily be seen) to the past of the bump, this same solution will be identifiable with the (partly positive-frequency, partly negative-frequency) Minkowski-space Klein-Gordon solution

$$\phi_{\mathrm{in}}(t, \mathbf{x}) = \left((2\mu)^{-1/2} \exp(-i\mu t\right) \alpha a)(\mathbf{x}) + \left((2\mu)^{-1/2} \exp(i\mu t)\bar{\beta}a\right)(\mathbf{x})$$

and this could be taken to be the defining equation for the operators α and β.

It is then known (by a 1962 theorem of Shale) that the automorphism (5) will be unitarily implemented if and only if β is a Hilbert Schmidt operator on $\mathcal{H}$. Wald (1979, in case $m \geq 0$) and Dimock (1979, in case $m \neq 0$) have verified that this condition is satisfied in the case of our bump-of-curvature situation. In that case, if we denote the unitary implementor by U, we have the results

(i) The expectation value $\langle U\Omega | N(a) U\Omega \rangle_{\mathcal{F}(\mathcal{H})}$ of the number operator, $N(a) = \hat{a}^\dagger(a)\hat{a}(a)$, where a is a normalized element of $\mathcal{H}$, is equal to $\langle \beta a | \beta a \rangle_{\mathcal{H}}$.

(ii) First note that there exists an orthonormal basis of vectors, e_i, ($i = 1 \ldots \infty$), in $\mathcal{H}$ such that the (Hilbert Schmidt) operator $\bar{\beta}^* \bar{\alpha}^{*-1}$ has the canonical form $\sum_i \lambda_i \langle Ce_i | \cdot \rangle | e_i \rangle$. We then have (up to an undetermined phase)

$$U\Omega = N \exp\left(-\frac{1}{2} \sum_i \lambda_i \hat{a}^\dagger(e_i) \hat{a}^\dagger(e_i) \right) \Omega.$$

where the normalization constant N is chosen so that $\|U\Omega\| = 1$ This formula makes manifest that the particles are created in pairs.

We remark that, identifying elements, a, of $\mathcal{H}$ with positive-frequency solutions (below, we shall call them "modes") as explained above, Result (i) may alternatively be expressed by saying that the expectation value, $\omega_{\mathrm{in}}(N(\mathrm{a}))$, in the *in-vacuum state* of the occupation number, $N(\mathrm{a})$, of a *normalized mode*, a, to the future of the bump is given by $\langle \beta a | \beta a \rangle_{\mathcal{H}}$.

This formalism and the results, (i) and (ii) above, will generalize (at least heuristically, and sometimes rigorously – see especially the rigorous scattering theoretic work in the 1980's by Dimock and Kay and more recently by A. Bachelot and others) to more realistic spacetimes which are only asymptotically flat or asymptotically stationary. In favourable cases, one will still have notions of classical solutions which are positive-frequency asymptotically towards the future/past, and, in consequence, one will have well-defined asymptotic notions of "vacuum" and "particles". Also, in, e.g. cosmological, models where the background spacetime is slowly-varying in time, one can define approximate *adiabatic* notions of classical positive frequency solutions, and hence also of quantum "vacuum" and "particles" at each finite value of the cosmological time. But, at times where the gravitational field is rapidly varying, one does not expect there to be any sensible notion of "particles". And, in a rapidly time-varying background gravitational field which never settles down one does not expect there to be any sensible particle interpretation of the theory at all. To understand these statements, it suffices to consider the $1+0$-dimensional Klein-Gordon equation with an external potential V:

$$\left(-\frac{d^2}{dt^2} - m^2 - V(t)\right)\phi = 0$$

which is of course a system of one degree of freedom, mathematically equivalent to the harmonic oscillator with a time-varying angular frequency $\varpi(t) = (m^2+V(t))^{1/2}$. One could of course express its quantum theory in terms of a time-evolving Schrödinger wave function $\Psi(\varphi, t)$ and attempt to give this a particle interpretation at each time, s, by expanding $\Psi(\varphi, s)$ in terms of the harmonic oscillator wave functions for a harmonic oscillator with some particular choice of angular frequency. But the problem is, as is easy to convince oneself, that there is no such good choice. For example, one might think that a good choice would be to take, at time s, the set of harmonic oscillator wave functions with angular frequency $\varpi(s)$. (This is sometimes known as the method of "instantaneous diagonalization of the Hamiltonian"). But suppose we were to apply this prescription to the case of a smooth $V(\cdot)$ which is constant in time until time 0 and assume the initial state is the usual vacuum state. Then at some positive time s, the number of particles predicted to be present is the same as the number of particles predicted to be present on the same prescription at all times after s for a $\hat{V}(\cdot)$ which is equal to $V(\cdot)$ up to time s and then takes the constant value $V(s)$ for all later times (see Figure 2). But $\hat{V}(\cdot)$ will generically have a sharp corner in its graph (i.e. a discontinuity in its time-derivative) at time s and one would expect a large part of the particle production in the latter situation to be accounted for by the presence of this sharp corner – and therefore a large part of the predicted particle-production in the case of $V(\cdot)$ to be spurious.



Figure 2 Plots of ϖ against t for the two potentials V (continuous line) and $\hat{V}$ (continuous line up to s and then dashed line) which play a role in our critique of "instantaneous diagonalization".

Back in 1+3 dimensions, even where a good notion of particles is possible, it depends on the choice of time-evolution, as is dramatically illustrated by the Unruh effect (see Section 5).

4 Theory of the Stress-Energy Tensor

To orient ideas, consider first the free (minimally coupled) scalar field, $(\Box - m^2)\phi = 0$, in Minkowski space. If one quantizes this system in the usual Minkowski-vacuum representation, then the expectation value of the *renormalized stress-energy tensor* (which in this case is the same thing as the *normal ordered stress-energy tensor*) in a vector state Ψ in the Fock space will be given by the formal *point-splitting* expression

$$\langle\Psi|T_{ab}(x)\Psi\rangle = \lim_{(x_1,x_2)\to(x,x)} \left(\partial_a^1\partial_b^2 - \frac{1}{2}\eta_{ab}(\eta^{cd}\partial_c^1\partial_d^2 + m^2)\right) \left(\langle\Psi|\rho_0(\phi(x_1)\phi(x_2))\Psi\rangle - \langle\Omega^{\mathcal{F}}|\rho_0(\phi(x_1)\phi(x_2))\Omega^{\mathcal{F}}\rangle\right) \tag{6}$$

where η_{ab} is the usual Minkowski metric. A sufficient condition for the limit here to be finite and well-defined would, e.g., be for Ψ to consist of a (normalised) finite superposition of n-particle vectors of form $\hat{a}^\dagger(a_1), \ldots, \hat{a}^\dagger(a_n)\Omega^{\mathcal{F}}$ where the smearing functions $a_1, \ldots, a_n$ are all C^∞ elements of $\mathcal{H}$ (i.e. of $L^2_{\mathbf{C}}(\mathbf{R}^3)$). The reason this works is that the two-point function in such states shares the same short-distance singularity as the Minkowski-vacuum two-point function. For exactly the same reason, one obtains a well-defined finite limit if one defines the expectation value of the stress-energy tensor in any physically admissible quasi-free state by the expression

$$\omega(T_{ab})(x)) = \lim_{(x_1,x_2)\to(x,x)} \left(\partial_a^1\partial_b^2 - \frac{1}{2}\eta_{ab}(\eta^{cd}\partial_c^1\partial_d^2 + m^2)\right) \left(\omega(\phi(x_1)\phi(x_2)) - \omega_0(\phi(x_1)\phi(x_2))\right). \tag{7}$$

This latter point-splitting formula generalizes to a definition for the *expectation value of the renormalized stress-energy tensor* for an arbitrary physically

admissible quasi-free state (or indeed for an arbitrary state whose two point function has *Hadamard form* – i.e. whose anticommutator function satisfies Condition (d) of Section 2) on the minimal field algebra and to other linear field theories (including the stress tensor for a conformally coupled linear scalar field) on a general globally hyperbolic spacetime (and the result obtained agrees with that obtained by other methods, including *dimensional regularization* and *zeta-function regularization*). However, the generalization to a curved spacetime involves a number of important new features which we now briefly list. (See (Wald, 1978) for details.)

First, the subtraction term which replaces $\omega_0(\phi(x_1)\phi(x_2))$ is, in general, not the expectation value of $\phi(x_1)\phi(x_2)$ in any particular state, but rather a particular *locally constructed Hadamard two-point function* whose physical interpretation is more subtle; the renormalization is thus in general not to be regarded as a normal ordering. Second, the immediate result of the resulting limiting process will not be covariantly conserved and, in order to obtain a covariantly conserved quantity, one needs to add a particular *local geometrical correction term*. The upshot of this is that the resulting expected stress-energy tensor is covariantly conserved but possesses a (state-independent) *anomalous trace*. In particular, for a massless conformally coupled linear scalar field, one has (for all physically admissible quasi-free states, ω) the *trace anomaly formula*

$$\omega(T^a_a(x)) = (2880\pi^2)^{-1}(C_{abcd}C^{abcd} + R_{ab}R^{ab} - \frac{1}{3}R^2)$$

– plus an arbitrary multiple of $\Box R$. In fact, in general, the thus-defined renormalized stress-energy tensor operator (see below) is only defined up to a *finite renormalization ambiguity* which consists of the addition of arbitrary multiples of the functional derivatives with respect to g_{ab} of the four quantities

$$I_n = \int_{\mathcal{M}} F_n(x)|\det(g)|^{1/2}d^4x,$$

$n = 1\ldots 4$, where $F_1 = 1$, $F_2 = R$, $F_3 = R^2$, $F_4 = R_{ab}R^{ab}$. In the Minkowski-space case, only the first of these ambiguities arises and it is implicitly resolved in the formulae (6), (7) inasmuch as these effectively incorporate the *renormalization condition* that $\omega_0(T_{ab}) = 0$. (For the same reason, the locally-flat example we give below has no ambiguity.)

One expects, in both flat and curved cases, that, for test functions, $F \in C_0^\infty(\mathcal{M})$, there will exist operators $T_{ab}(F)$ which are *affiliated to* the net of local W^*-algebras referred to in Section 2 and that it is meaningful to write

$$\int_{\mathcal{M}} \omega(T_{ab}(x))F(x)|\det(g)|^{1/2}d^4x = \omega(T_{ab}(F))$$

provided that, by ω on the right-hand side, we understand the extension of ω from the Weyl algebra to this net. ($T_{ab}(F)$ is however not expected to

belong to the minimal algebra or be affiliated to the Weyl algebra.)

An interesting simple example of a renormalized stress-energy tensor calculation is the so-called *Casimir effect* calculation for a linear scalar field on a (for further simplicity, $1+1$-dimensional) *timelike cylinder spacetime* of radius R (see Figure 3). This spacetime is globally hyperbolic and stationary and, while locally flat, globally distinct from Minkowski space. As a result, while – provided the regions $\mathcal{O}$ are sufficiently small (such as the diamond region in Figure 3) – elements $\mathcal{A}(\mathcal{O})$ of the minimal net of local algebras on this spacetime will be identifiable, in an obvious way, with elements of the minimal net of local algebras on Minkowski space, the stationary ground state ω_{cylinder} will, when restricted to such thus-identified regions, be distinct from the Minkowski vacuum state ω_0. The resulting renormalized stress-energy tensor (as first pointed out in (Kay, 1979), defineable, once the above identification has been made, exactly as in (7)) turns out to be non-zero and, interestingly, to have a (in the natural coordinates, constant) *negative energy density* T_{00}. In fact:

$$\omega_{\text{cylinder}}(T_{ab}) = \frac{1}{24\pi R^2}\eta_{ab}.$$



Figure 3 The time-like cylinder spacetime of radius R with a diamond region isometric to a piece of Minkowski space. See (Kay, 1979).

5 Hawking and Unruh effects

The original (1974) calculation by Hawking concerned a model spacetime for a star which collapses to a black hole. For simplicity, we shall only discuss the spherically-symmetric case. (See Figure 4.)



Figure 4 The spacetime of a star collapsing to a spherical black hole.

Adopting a similar "mode" viewpoint to that mentioned after Results (i) and (ii) in Section 3, the result of the calculation may be stated as follows: For a real linear scalar field satisfying (2) with $m = 0$ (and $V = 0$) on this spacetime, the expectation value $\omega_{\rm in}(N(a_{\varpi,\ell}))$ of the occupation number of a one-particle outgoing mode $a_{\varpi,\ell}$) localized (as far as a normalized mode can be) around ϖ in angular-frequency-space and about retarded time v and with angular momentum "quantum number" ℓ, in the in-vacuum state (i.e. on the minimal algebra for a real scalar field on this model spacetime) $\omega_{\rm in}$ is, at late retarded times, given by the formula

$$\omega_{\rm in}(N(a_{\varpi,\ell})) = \frac{\Gamma(\varpi,\ell)}{\exp(8\pi M\varpi) - 1}$$

where M is the mass of the black hole and the *absorption factor* (alternatively known as *grey body factor*) $\Gamma(\varpi,\ell)$ is equal to the norm-squared of that part of the one-particle mode, $a_{\varpi,\ell}$, which, viewed as a complex positive frequency classical solution propagating backwards in time from late retarded times, would be absorbed by the black hole. This calculation can be understood as an application of Result (i) of Section 3 (even though the spacetime is more complicated than one with a localized "bump of curvature" and even though the relevant overall time-evolution will not be unitarily implemented, the result still applies when suitably interpreted) and the heart of the calculation is an asymptotic estimate of the relevant "β" Bogoliubov coefficient which turns out to be dependent on the geometrical optics of rays which pass through the star just before the formation of the horizon. This result suggests that the in-vacuum state is indistinguishable at late retarded times from a state of black-body radiation at the *Hawking temperature*, $T_{\rm Hawking} = 1/8\pi M$, in Minkowski space from a black-body with the same absorption factor. This was confirmed by further work by many authors. Much of that work, as well as the original result of Hawking was partially heuristic but later work by Dimock and Kay (1987), by

Fredenhagen and Haag (1990) and by Bachelot (1999) and others has put different aspects of it on a rigorous mathematical footing. The result generalizes to non-zero mass and higher spin fields and to interacting fields as well as to other types of black hole and the formula for the Hawking temperature generalizes to

$$T_{\text{Hawking}} = \kappa/2\pi$$

where κ is the *surface gravity* of the black hole.

This result suggests that there is something fundamentally "thermal" about quantum fields on black-hole backgrounds and this is confirmed by a number of mathematical results. In particular, the theorems in the two papers (Kay and Wald, 1991) and (Kay, 1993), combined together, tell us that there is a unique state on the Weyl algebra for the *maximally extended Schwarzschild spacetime* (a.k.a. *Kruskal-Szekeres spacetime*) (see Figure 5) which is invariant under the *Schwarzschild isometry group* and whose two-point function has Hadamard form. Moreover, they tell us that this state, when restricted to a single wedge (i.e. the exterior Schwarzschild spacetime) is necessarily a KMS state at the Hawking temperature.



Figure 5 The geometry of maximally extended Schwarzschild (/or Minkowski) spacetime. In the Schwarzschild case, every point represents a two-sphere (/in the Minkowski case, a two-plane). The curves with arrows on them indicate the Schwarzschild time-evolution (/one-parameter family of Lorentz boosts). These curves include the (straight lines at right angles) event horizons (/Killing horizons).

This unique state is known as the *Hartle-Hawking-Israel state*. These results in fact apply more generally to a wide class of globally hyperbolic spacetimes with *bifurcate Killing horizons* including de Sitter space – where the unique state is sometimes called the *Euclidean* and sometimes the *Bunch-Davies* vacuum state – as well as to Minkowski space, in which case the

unique state is the usual Minkowski vacuum state, the analogue of the exterior Schwarzschild wedge is a so-called *Rindler wedge*, and the relevant isometry group is a one-parameter family of wedge-preserving Lorentz boosts. In the latter situation, the fact that the Minkowski vacuum state is a KMS state (at "temperature" $1/2\pi$) when restricted to a Rindler wedge and regarded with respect to the time-evolution consisting of the wedge-preserving one-parameter family of Lorentz boosts is known as the *Unruh effect* (1975). This latter property of the Minkowski vacuum in fact generalizes to general *Wightman quantum field theories* and is in fact an immediate consequence of a combination of the *Reeh Schlieder Theorem* (applied to a Rindler wedge) and the *Bisognano Wichmann Theorem* (1975). The latter theorem says that the defining relation (1) of a KMS state holds if, in (1), we identify the operator J with the complex conjugation which implements wedge reflection and H with the self-adjoint generator of the unitary implementor of Lorentz boosts. We remark that the Unruh effect illustrates how the concept of "vacuum" (when meaningful at all) is dependent on the choice of time-evolution under consideration. Thus the usual Minkowski vacuum is a ground state with respect to the usual Minkowski time-evolution but not (when restricted to a Rindler wedge) with respect to a one-parameter family of Lorentz boosts; with respect to these, it is, instead, a KMS state.

6 Non-Globally Hyperbolic Spacetimes and the "Time Machine" Question

In (Hawking, 1992) it is argued that a spacetime in which a time-machine gets manufactured should be modelled (see Figure 6) by a spacetime with an *initial globally hyperbolic region* with a region containing *closed timelike curves* to its future and such that the future boundary of the globally hyperbolic region is a *compactly generated Cauchy horizon*. On such a spacetime, (Kay, Radzikowski and Wald, 1997) proves that it is impossible for any distributional bisolution which satisfies (even a certain weakened version of) the Hadamard condition on the initial globally hyperbolic region to continue to satisfy that condition on the full spacetime – the (weakened) Hadamard condition being necessarily violated at at least one point of the Cauchy horizon. This result implies that, however one extends a state, satisfying our conditions (a), (b), (c) and (d), on the minimal algebra for (2) on the initial globally hyperbolic region, the expectation value of its stress-energy tensor must necessarily become singular *on* the Cauchy horizon. This result, together with many heuristic results and specific examples considered by many other authors appears to support the validity of the (Hawking, 1992) *chronology protection conjecture* to the effect that it is impossible in principle to manufacture a time machine. However, there are potential loopholes in the physical interpretation of this result as pointed out by Visser (1997), as

well as other claims by various authors that one can nevertheless violate the chronology protection conjecture. For a recent discussion on this question, we refer to (Visser, 2003).



Figure 6 The schematic geometry of a spacetime in which a time-machine gets manufactured.

7 Other Related Topics and Some Warnings

There is a vast computational literature, calculating the expectation values of stress-energy tensors in states of interest for scalar and higher spin linear fields (and also some work for interacting fields) on interesting cosmological and black-hole backgrounds. Quantum field theory on de Sitter and anti-de Sitter space is a big subject area in its own right with recent renewed interest because of its relevance to *string theory* and *holography*. Also important on black hole backgrounds is the calculation of grey-body factors, again with renewed interest because of relevance to string theory and to *brane-world* scenarios.

There are many further mathematically rigorous results on algebraic and axiomatic quantum field theory in a curved spacetime setting, including versions of *PCT*, *Spin-Statistics* and *Reeh-Schlieder* theorems and also rigorous *energy inequalities* bounding the extent to which expected energy densities can be negative etc.

There is much mathematical work controlling scattering theory on black holes, partly with a view to further elucidating the Hawking effect.

Perturbative renormalization theory of interacting quantum fields in curved spacetime is also now a highly developed subject.

Beyond quantum field theory in a fixed curved spacetime is *semiclassical gravity* which takes into account the *back reaction* of the expectation value of the stress-energy tensor on the classical gravitational background. There are also interesting condensed-matter analogues of the Hawking effect such as *dumb holes*.

Readers exploring the wider literature, or doing further research on the

subject should be aware that the word "vacuum" is sometimes used to mean "ground state" and sometimes just to mean "quasi-free state". Furthermore they should be cautious of attempts to define particles on Cauchy surfaces in *instantaneous diagonalization* schemes (cf. the remarks at the end of Section 3). When studying (or performing) calculations of the "expectation value of the stress-energy tensor" it is always important to ask oneself with respect to which state the expectation value is being taken. It is also important to remember to check that candidate two-point (anticommutator) functions satisfy the positivity condition (c) of Section 2. Typically two-point distributions obtained via *mode sums* automatically satisfy Condition (c) (and Condition (d)), but those obtained via *image* methods don't always satisfy it. (When they don't, the presence of non-local spacelike singularities is often a tell-tale sign as can be inferred from Kay's Conjecture/Radzikowski's Theorem discussed in Section 2.) There are a number of apparent implicit assertions in the literature that some such two-point functions arise from "states" when of course they can't. Some of these concern proposed analogues to the Hartle-Hawking-Israel state for the (appropriate maximal globally hyperbolic portion of the maximally extended) Kerr spacetime. That they can't belong to states is clear from a theorem in (Kay and Wald, 1991) which states that there is no stationary Hadamard state on this spacetime at all. Others of them concern claimed "states" on spacetimes such as those discussed in Section 6 which, if they really were states would seem to be in conflict with the chronology protection conjecture. Finally, beware states (such as the so-called *α-vacua* of de Sitter spacetime) whose two-point distributions violate the "Hadamard" Condition (d) of Section 2 and which therefore do not have a well-defined finite expectation value for the renormalized stress-energy tensor.

8 See also

Black Hole Thermodynamics. **Algebraic Approach to Quantum Field Theory**.

9 Further Reading

DeWitt BS (1975) Quantum field theory in curved space-time. *Physics Reports* 19, No. 6, 295-357.

Birrell ND and Davies PCW (1982) *Quantum Fields in Curved Space.* Cambridge: Cambridge University Press.

Haag R (1996) *Local Quantum Physics.* Berlin: Springer.

Dimock J (1980) Algebras of local observables on a manifold. *Communications in Mathematical Physics* 77: 219-228.

Brunetti R, Fredenhagen K and Verch R (2003) The generally covariant locality principle – A new paradigm for local quantum physics. *Communications in Mathematical Physics* 237:31-68.

Kay, BS (2000) Application of linear hyperbolic PDE to linear quantum fields in curved spacetimes: especially black holes, time machines and a new semi-local vacuum concept. Journées *Équations aux Dérivées Partielles*, Nantes, 5-9 juin 2000 GDR 1151 (CNRS): IX1-IX19. (Also available at http://www.math.sciences.univ-nantes.fr/edpa/2000/html or as gr-qc/0103056.)

Wald RM (1978) Trace anomaly of a conformally invariant quantum field in a curved spacetime. *Physical Review* D17: 1477-1484.

Kay BS (1979) Casimir effect in quantum field theory. (*Original title:* The Casimir effect without magic.) *Physical Review* D20: 3052-3062.

Hawking SW (1975) Particle Creation by Black Holes. *Communications in Mathematical Physics* 43: 199-220.

Kay BS and Wald RM (1991) Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on spacetimes with a bifurcate Killing horizon. *Physics Reports* 207, No. 2, 49-136.

Kay BS (1993) Sufficient conditions for quasifree states and an improved uniqueness theorem for quantum fields on space-times with horizons. *Journal of Mathematical Physics* 34: 4519-4539.

Wald RM (1994) *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics.* Chicago: University of Chicago Press.

Hartle JB and Hawking SW (1976) Path-integral derivation of black-hole radiance. *Physical Review* D13: 2188-2203.

Israel W (1976) Thermo-field dynamics of black holes. *Physics Letters A* 57: 107-110.

Unruh W (1976) Notes on black hole evaporation. *Physical Review* D14: 870-892.

Hawking SW (1992) The Chronology Protection Conjecture. *Physical Review* D46: 603-611.

Kay BS, Radzikowski MJ and Wald RM (1997) Quantum field theory on spacetimes with a compactly generated Cauchy horizon. *Communications in Mathematical Physics* 183: 533-556.

Visser M (2003) The quantum physics of chronology protection. In: Gibbons GW, Shellard EPS and Rankin SJ *The Future of Theoretical Physics and Cosmology.* Cambridge: Cambridge University Press.